

# Hyperhomocysteinemia, Diabetes and Cardiovascular Disease

THE HOORN STUDY



## Contents

Chapter 1	General introduction	9
Chapter 2	Hyperhomocysteinemia is associated with an increased risk of cardiovascular disease, especially in non-insulin-dependent diabetes mellitus: A population-based study. <i>Arterioscler Thromb Vasc Biol</i> 1998;18:133-138	29
Chapter 3	Hyperhomocysteinemia is associated with aortoiliac but not with isolated crural arterial occlusive disease: The Hoorn Study.	45
Chapter 4	Hyperhomocysteinemia increases risk of 5-year mortality, especially in type 2 diabetes: The Hoorn Study.	59
Chapter 5	Serum homocysteine level and protein intake are related to risk of microalbuminuria: The Hoorn Study. <i>Kidney Int</i> 1998;54:203-209	75
Chapter 6	Hyperhomocysteinemia is related to risk of retinopathy in type 2 diabetes: The Hoorn Study.	93
Chapter 7	Hyperhomocysteinemia is not related to risk of distal somatic polyneuropathy: The Hoorn Study.	109
Chapter 8	Does metformin increase the serum total homocysteine level in non-insulin-dependent diabetes mellitus? <i>J Intern Med</i> 1997;242:389-394	121
Chapter 9	General discussion	133
	Abbreviations	145
	Summary	147
	Samenvatting	151
	Dankwoord	155
	About the author	157